

**UNCLASSIFIED**

---

**AD 400 512**

*Reproduced  
by the*

**ARMED SERVICES TECHNICAL INFORMATION AGENCY  
ARLINGTON HALL STATION  
ARLINGTON 12, VIRGINIA**



---

**UNCLASSIFIED**

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

63-3-1

FTD-TT- 63-162

CATALOGED BY ASTIA  
AS AD-NO-400 512


## TRANSLATION

THE "TOXIN-ENZYME" HYPOTHESIS AND THE ROLE OF BACTERIAL  
ENZYMES IN THE PATHOGENESIS OF INFECTIONS

By

I. V. Domaradskiy

## FOREIGN TECHNOLOGY DIVISION



AIR FORCE SYSTEMS COMMAND

WRIGHT-PATTERSON AIR FORCE BASE  
OHIO

## UNEDITED ROUGH DRAFT TRANSLATION

THE "TOXIN-ENZYME" HYPOTHESIS AND THE ROLE OF  
BACTERIAL ENZYMES IN THE PATHOGENESIS OF  
INFECTIONS

By: I. V. Domaradskiy

English Pages: 24

SOURCE: Russian Periodical, Uspekhi Sovremennoy  
Biologii, Vol. 54, Nr. 1, 1962, pp. 57-70.

T-19  
SOV/221-62-54-1-1/1

THIS TRANSLATION IS A RENDITION OF THE ORIGINAL FOREIGN TEXT WITHOUT ANY ANALYTICAL OR EDITORIAL COMMENT. STATEMENTS OR THEORIES ADVOCATED OR IMPLIED ARE THOSE OF THE SOURCE AND DO NOT NECESSARILY REFLECT THE POSITION OR OPINION OF THE FOREIGN TECHNOLOGY DIVISION.

PREPARED BY:

TRANSLATION SERVICES BRANCH  
FOREIGN TECHNOLOGY DIVISION  
WP-AFB, OHIO.

FTD-TT-63-162/1+4

Date 12 Mar 1963

THE "TOXIN-ENZYME" HYPOTHESIS AND THE ROLE OF  
BACTERIAL ENZYMES IN THE PATHOGENESIS OF INFECTIONS

I. V. Domaradskiy

In recent years many works have appeared in which the authors attempted to identify bacterial toxins\* with enzymes, and toxicosis is considered the consequence of the action of these enzymes on appropriate substrates. It is assumed that death of an organism ensues either as a result of the destruction of cell structures or under the effect of toxic products which are formed secondarily (Konikov, 1950; Gubarev, 1952, 1956; Galaev, 1956; Gubarev and Ivanovskiy, 1958; Matveyev, 1959; Zaplatina, 1959; Poverenny, 1961 et al.).

1.

As is known the "toxin-enzyme" hypothesis arose as a result of studying gram-positive bacteria. It is precisely in these bacteria that toxins having catalytic properties were found. And this is not by chance.

All species leading a parasitic mode of life are found vastly more rarely among gram-positive bacteria than in the group of

\* In this article we will deal mainly with the so-called exotoxins.

gram-negative bacteria. Many of them, especially the sporiferous (Bac. botulinus, Bac. tetani, Bac. anthracis, Bac. sordellii, Bac. perfringens and other causative agents of gas gangrene) are frequent (or permanent) inhabitants of the external environment, including the soil (Mishustin and Pertsovskaya, 1954; Matveyev, 1959, 1960). In view of this the collection of enzymes of pathogenic gram-positive bacteria is distinguished by a great wealth and diversity. In particular they are widely represented by hydrolases, enzymes which accomplish the first steps in the breakdown of complex organic substances which cannot be per se assimilated by cells. These enzymes are apparently excreted into the ambient medium. Other enzymes participating in intermediate metabolisms are usually not excreted to the external environment or pass to it only after cell necrosis. Those that are called toxins are also excreted to the external environment and are among the gram-positive bacteria. Pathogenic gram-negative bacteria, with the exception of Bact. dysenteriae Shiga, which by virtue of their paracitic mode of life mainly have intracellular enzymes, do not produce true toxins.

Finally, judging by the data in the literature, all toxins or factors of aggression having catalytic properties are hydrolases.\*

Therefore, there is nothing surprising that toxins-enzymes are found only in representatives of gram-positive microbes and only in those whose toxin production is directly associated with processes of cell nutrition.

From our point of view it would be rash to apply the data on the identity of enzymes with toxins obtained in the study of this narrow group of bacteria to all microorganisms.

---

\* The exception is only the water-soluble oxidase of 1-amino acid found in snake venom (Zeller, Maritz, 1944).

A great similarity does exist between toxins and enzymes and this is noted when there were still no clear concepts of the chemical nature of the latter (Wells, 1929). One of the features of this similarity is the toxicity inherent to certain enzymes upon parenteral administration. A classical example confirming this is crystalline urease which causes breakdown of blood urea and, as a consequence, of this ammonia poisoning of the organism (Kirk, Sumner, 1931); birds whose urea is not the final product of nitrogen metabolism, are resistant to urease (Howell, 1932). Another example is the toxicity of crystalline trypsin and chymotrypsin for rabbits, rats, guinea pigs, cats, and dogs (Rocha e Silva, 1931a, b; 1940). Both these enzymes cause myocardial damage to animals (Hendley et al., 1956). Cathepsin, formed by streptococci of group A, has a similar effect on rabbit myocardium (Kellner, Robertson, 1954a, b). It is possible to give other examples indicating the toxicity of enzymes.

The features of similarity between enzymes and toxins also include their ability to cause the formation of corresponding antibodies. True, enzymes are rather weak antigens (Dormarabskiy, 1961).

Under the effect of formaldehyde and heating, certain enzymes can be changed to anaenzymes. This is proved by the example of papain and gelatinase of the tetanus bacillus (Ramon, 1957) which under the indicated effects lose their enzymic activity but retain the antigen properties. Serum of animals immunized with "anapapain" or "anagelatinase" has the ability to suppress the activity of the corresponding enzymes not treated with formalin and not subjected to heating. We can conclude from this that the active centers of enzymes are apparently not identical to the determinative groups causing antigen properties. The similarity of enzymes with toxins is also manifested in this.

Minally, enzymes like toxins are proteins; most of them are thermostable, although in certain cases they can withstand the effect of high temperatures without a decrease in specific activities (Thompson et al., 1958).

However, the similarities between toxins and enzymes is also insufficient for their identity since there are no indisputable proofs of the presence of catalytic properties for the overwhelming majority of toxins.

In spite of all this there is a tendency to enlarge the frameworks of the "toxin-enzyme" hypothesis and to convert it to a universal theory of the action of toxins. The presence of catalytic properties is even postulated for such toxins as diphtheria, botulinus and tetanus (Gubarev, 1952). The facts testifying against this hypothesis are not sufficiently taken into account.

The latent period in the action of toxins and their unusual activity force us to assume that they act not like enzymes but like compounds suppressing the transformation and synthesis of certain of the most important cell components (Dubos, 1957). In particular, the available data indicate that diphtheria toxin in sensitive animals block the synthesis of cytochrome b or one of the enzymes close to it (Pappenheimer, Hendee, 1947; Pappenheimer, Williams, 1952; Pappenheimer, 1947a, b). Moreover diphtheria toxin noticeably inhibits the synthesis of protein of human carcinoma. In this respect its action differs from the action of cyanide and dinitrophenol (Strauss, Hendee, 1959). Another toxin produced by the fungus *Fusarium lycopersici* and causing wilting and leafroll of tomatoes affects metabolism because by its structure it is very similar to the growth factor streptogenin. In this case between the toxin, which is a peptide derivative of asparagin, glycine, and serine, and streptogenin whose composition includes



glutamic acid and glycine, there is an antagonism of a noncompetitive nature (for details see Wooley, 1954).

The so-called endotoxins of gram-negative bacteria, which in most cases are lipo-polysacchride-protein complexes completely devoid of any catalytic properties, evidently act according to the type of inhibitors. A characteristic feature of the action of endotoxins is that regardless of their origin they cause a similar picture of intoxication and similar biochemical shifts in an organism (Zil'ber, 1958; Dubos, 1957; Domaradskiy et al., 1961; Delaunay, Lebrun, 1947 et al.). According to the opinion of Heyningen (1950), this feature should underlie the differentiation of toxins of gram-negative and gram-positive bacteria since each toxin of the latter has its own specificity of action.

Unfortunately we still do not know the points of application of the action of most toxins. We can only say that in the case of the toxins of the causative agents of tetanus and botulism, cholinesterase of the main ganglia of the central nervous system is not such a point (Isibasi, 1959).\*

The instability of the condition of the adherents of the "toxin-enzyme" hypothesis, from our point of view, is easily demonstrated by the following examples.

First, the investigators most frequently have to deal with unpurified toxic fractions of microbes in which other biologically active substances are present together with the toxins proper. Many of

\* Buehler et al. (1947) considers that when explaining the high toxicity of botulinus toxin it is necessary to take into account: 1) the polar nature of the compound, 2) the arrangement of amino acids within the molecule and the geometric configuration of protein as a whole, 3) the possibility of an atypical peptide bond between the aspartic and glutamic acids, 4) the possibility of the presence of toxic groups in unanalyzable amounts, 5) the high molecular weight of protein.

them have a catalytic activity. For example staphylococcal enterotoxin preparation contained apyrase wherein it is still impossible to determine where the action of enterotoxin ends and the action of apyrase begins (Sugiyama, Dack, 1955). The same can be said about o-streptolysin whose leukotoxicity and ability to inhibit respiration of mitochondria are associated with the presence of diphosphopyridine nucleotide (Carlson et al., 1956, 1957), or about staphylocoagulases containing lipase (Drummond, Tager, 1959a, b), etc. An underestimation of this circumstance can lead to serious errors.

Second, it is difficult to explain why one and the same enzymes excreted by various microbes sharply differ in their toxicity. For example, little understood is the reason for the approximately similar toxicity of the lecithinases of the pathogenic microbe *Bac. perfringens* and the nonpathogenic microbe *Bac. cereus* in spite of their dissimilarities as antigens (Chu, 1949) and the absence of toxicity for lecithinase of the anthrax causative agent (Costlow, 1958). An explanation of the relatively lesser toxicity of the lecithinase of *Bac. cereus* in comparison with that of the lecithinase of *Bac. perfringens* is that the former is suppressed by normal serum which apparently is not applicable to lecithinase of any origin since the nontoxic lecithinase of *Bac. bifementans* is not suppressed by normal serum (Miles, Miles, 1950).

Third, since it was established that the participation of active centers of an enzyme in the actions of an enzyme with an antibody is not obligatory (Kirk, 1933; Kirk, Sumner, 1934; Haas, 1940; Campbell, Fourt, 1939), we cannot yet give an answer to two questions: 1) does the toxicity of enzymes depend only on their active group and 2) what explains the mechanism of the therapeutic actions of corresponding antiserums.

Fourth, acceptance that enzymes have a direct toxic action on tissue, cells, or metabolism does not explain (or does not always explain) the cause of their over-all action on an organism. It is not precluded that in addition to enzymic action, toxins-enzymes have other properties no less important for the origination of disease (Dubos, 1957; Heyning, 1958).

Finally, different enzymes and substances not having a catalytic activity cannot cause similar symptoms of a disease or a similar reaction with respect to isolated organs. For example, the permeability of capillaries is increased not only by hyaluronidase, but also by trypsin, chymotrypsin, plasmin and proteinase of the skin (Paskhina, 1955). Erythrocytes are dissolved by saponins and also by the  $\alpha$ -toxin of *Bac. perfringens* and the proteinase of *staphylococcus* (Robinson et al., 1960). A picture of beri-beri can be caused by  $B_1$  avitaminosis of an exogenous origin, by infection of the organ with *Bac. thiaminolyticus* and by protein deficiency. Common symptoms are noted in tetanus and strychnism.

In all these cases (as in many others) we deal with the same consequence but with a different mechanism of action.

It follows from what has been stated that when solving the problem of the mode of action of toxins we must show extreme care, and a particular conclusion obtained on the basis of studying the toxins of individual species of bacteria cannot unconditionally be applied to other toxins. The method of analysis, no matter how attractive it seems, is not applicable in this case.

## 2.

However, while we object to those who consider all toxins as enzymes we do not deny the role of enzymes of bacteria in the pathogenesis

the corresponding disease in some cases. Conversely, we consider that too little attention has been devoted to this aspect of the problem.

The ability to become adapted in a susceptible host, which is an "external environment" for a pathogenic microbe, to a considerable extent depends on the biochemical activity of the microbe. In contrast to microbiologists who judge the biochemical activity only by the behavior of bacteria on "variegated" series, we enclose a broader meaning in this concept. Biological activity is the intensity of metabolism, the intensity of all reactions occurring in a cell and causing its vital activity. From the indicated position not a single pathogenic microbe can be considered as biochemically inert. We will recall that the evolution of microbes mainly concern those processes which pertain to energy metabolism (fermentation, oxidation) and to a lesser extent affect anabolism. Anabolism of various organisms, including biochemically "active" or "inert" (from the point of view of microbiologists) faces one problem, "the building of body substances-protoplasts-, which in different representatives of life, according to present-day concepts, differ only in a slight degree" (Shaposhnikov, 1960).

A disease can be caused only by that pathogenic microbe which, all other things being equal (susceptibility of the macroorganism, state of the histohematic barriers, etc.), has a high biochemical activity sufficient to provide its source requirements of nutrition and energy. The deciding role in the development and outcome of a disease therefore belong to these bacterial enzymes.

There are many examples that bacterial enzymes can play the role of the pathogenic agents. One of them is the participation of thiaminase of *Bac. thiaminolyticus* in the development of B<sub>1</sub> avitaminosis

(Kleeberg, 1958). Another example is urease of *Proteus mirabilis* which causes the development of experimental pyelonephritis. In contrast to *Proteus*, *Escherichia coli*, deprived of urease activity, does not cause lesions of the renal epithelium either in vivo or in vitro (Braude, Sieminski, 1960).

The pathogenic significance of staphylococcal enzymes does not evoke doubts. Among them a particular place belongs to coagulase (Smith, Johnstone, 1958; Blobel, Berman, 1961), which, in the opinion of most investigators, is an enzyme (Lieb, 1960; Fusillo, Jaffurs, 1955). Fusillo and Jaffurs, (1955) consider it an adaptive enzyme.

Coagulase has antigen properties (Duthie, Lorenz, 1952; Blobel et al., 1960); animals actively immunized with coagulase are more resistant to infection by coagulase-positive strains of staphylococci (Boake, 1956).

Less is known about the pathogenic role of other enzymes in toxic filtrates of staphylococci. We can only note that lipase is inactivated by human blood gamma-globulins (Richou et al., 1960, 1961) and has no relation to blood clotting by coagulase (Drummond, Tager, 1959b) whereas protease, purified to a considerable extent of concomitant substances, has properties of  $\alpha$ -hemolysin (Robinson et al., 1960).

However, most typical is the example of the participation of the enzymes of *Bac. perfringens* and other bacteria in the pathogenesis of gas gangrene (for a detailed review see Kaplanskiy, 1954 and Chistovich, 1951). This was first proved by Macfarlane and Knight (1941) who identify  $\alpha$ -toxin of *Bac. perfringens* with the enzyme lecithinase (lecithinase C). The result of the action of this enzyme is the destruction of the protective layer of lecithin on particles of protein and fat and the destruction of cell structures, in particular

decomposition of the membranes of erythrocytes (Macfarlane, Oakley Anderson, 1941).

Lecithinase activity of  $\alpha$ -toxin is very great. For example, a single lethal dose of it can split all lecithin contained in mouse blood. The activity of lecithinase C is paralyzed by a specific antitoxin wherein competition for lecithinase exists between the antitoxin and lecithin (Zamecnik, Lipmann, 1947).

Macfarlane and MacLenan (1945), Oakley et al., (1946) and others detected another enzyme, collagenase, which dissolves collagen of connective tissue, in the composition of the toxin of *Bac. perfringens*.\* This enzyme, or kappa-toxin, is also neutralized by a specific antitoxin. In addition to collagenase, the filtrates of *Bac. perfringens* cultures contain other proteolytic enzymes, in particular lambda-enzyme. The latter was originally detected in a filtrate of a culture of *Bac. perfringens* type B and, according to the data of Oakley (1948), it can split proteins obtained from the skin of animals and also native collagen but does not act on animal muscles and denatured collagen. The problem as to what proteins are decomposed by collagenase and what ones by other proteolytic enzymes, including the lambda-enzyme, is in need of further investigation.

Nothing is yet known about the role of deoxyribonuclease of *Bac. perfringens* (Robb-Smith, 1945) in the pathogenesis of gas gangrene.

Among the bacterial enzymes we can presently consider only the lecithinase of certain clostridia and possibly the proteinase of staphylococcus as toxins (Robinson et al., 1960). Other enzymes in the sense in which the word "toxin" is usually used, are not toxins

---

\* It was recently established that collagenase of another causative agent of gas gangrene, *Bac. histolyticus*, has together with a proteolytic activity a trans-cis-propyl (hydroxyproline)-isomerase activity (Shpikiter, 1961).

because there is no specificity in their action on an organism, they do not kill tissue cells on which they act, and many of them (e.g., hyaluronidase, collagenase, deoxyribonuclease, certain fibrinolysins) decompose apparently exclusively extracellular substrates. These enzymes create only the necessary conditions for the development of microorganisms, the background, on which evolves the specific picture of the corresponding infectious process. Once again we repeat: this aspect of the activity of bacterial enzymes must not be underestimated.

In individual cases the changes in an organism caused by enzymes can be expressed very vigorously and in themselves serve as a cause of disease or prolong its course. It was established, for example, that the bacterial enzymes of dysentery caused the formation of different amines which in turn promote the formation of fatty degeneration of the liver in this infection (Takatori, 1955). The formation of amines and ammonia by intestinal bacteria also plays a role in the development of hepatic coma (Phear, Ruebner, 1956). Glutamine accumulates in brain tissue in pneumococcal infection of rats (Vyshepan and Porfir'eva, 1958). In the pathogenesis of cholera an important role is attributed to mucinase and lactic acid excreted by the cholera vib-  
0  
rion into the intestine. It is assumed that mucinase decomposes the epithelium and lactic acid increases the permeability of the intestines, thus fostering loss of liquids by the organism (Jenkin, Rowley, 1959). Enzymes of many bacteria decompose the group-specific substances of the blood (Iseki et al., 1959), cause intravascular coagulation (Smith, Johnstone, 1958; Blobel, Berman, 1961), intravital lysis and agglutination of arithrocytes (Keppie et al., 1955), cause pulmonary hemorrhages (Howes, Mandl, Zaffuto, 1960) and other phenomena.

3.

The question arises in connection with the role of enzymes in the pathogenesis of infection: what special features of the chemical composition or metabolism distinguish a pathogenic microbe from a non-pathogenic microbe?

As far as we know no substantial differences have been established in the chemical composition of pathogenic and nonpathogenic microbes.\*

The same amino acids are found, for example, in the proteins of pathogenic and nonpathogenic staphylococci (Sasaki, 1958). Moreover, the amino acid composition of the proteins of nonpathogenic cocci mainly coincides with that of the cholera and choleralike vibrios, differing from them only by the absence of an unidentified amino acid and aminobutyric acid. However, the spots of these amino acids are slightly demonstrated on certain chromatograms, which gives us grounds to consider this difference as being only quantitative (Samarina et al., 1950). The data on the identity of the amino acid composition of cholera and paracholera vibrios were confirmed by L. F. Pustovalova (1961).

True, many pathogenic and nonpathogenic microbes differ from one another with respect to the antigen structure, including with respect to polysacchrides. However, this is usually established by immunological and, very rarely, chemical methods. Proof of the relative ineffectiveness of serological methods of differentiation of pathogenic microbes from nonpathogenic is the fact of finding common antigens

---

\* As Tracey indicates (1954), "the chemical composition of the majority of organisms that have been studied is approximately the same. Glucose is the dominant sugar in potatoes and teredo, in the elephant and in bacteria. Many other sugars and about twenty amino acids within proteins are universal for all living organisms." A discussion of this question is in the works of Lise (1958) and Shaposhnikov (1960).



even in phylogenetically unrelated species of microorganisms, for example in pneumococcus and Friedlander's bacillus (Avery et al., 1925), in pneumococcus and various strains of yeasts (Neill, 1939).

It is even more difficult to establish the differences in the chemical composition of virulent and avirulent strains of pathogenic bacteria.

The  $V_1$  antigen is not an exclusive attribute of virulent strains of the typhoid bacillus, but is found in a number of other nonpathogenic species (Almon, 1943; Luippold, 1942, 1944; Stuart et al., 1943; Petrosyan and Zvenigorodskaya, 1957 etc.). Similarly the presence of M-protein in streptococci of group A does not always serve as an index of their virulence (Todd, Lancefield, 1928).

The virulent strains of *Escherichia coli* are similar to avirulent strains with respect to chemical composition, in particular the content of the toxic lipopolysaccharides (Harvey, Phillips, 1960). According to the data of Bennett (1959), the virulent strains of *Bac. anthracis* contain appreciably more phosphorous of DNA than the avirulent microorganism. The content of RNA was the same both for the virulent and avirulent strains. However, for three avirulent strains of the anthrax microbe sensitive to penicillin, the ratio of RNA and DNA was the same as for the virulent strains.

Apparently, a reliable difference in the chemical composition of virulent and avirulent strains could be established only for mycobacteria. In contrast to the avirulent strains, the virulent strains of mycobacteria contained a special lipid on the cell surface which causes their growth as coil-like bands. A single injection of this lipid (cord-factor), which per se does not cause morbid symptoms, accelerates the course of both acute and chronic tuberculous infection (cited from Model', 1958).

We can note further that not one of the autotrophic microorganisms, i.e., microbes which do not use organic substances as a primary energy source (Lise, 1958), has pathogenicity for man and animals. All pathogenic microbes belong to the number of heterotrophic organisms. However, the degree of heterotrophy of pathogenic microbes varies within wide limits. Some of them are parasites, do not grow at all or are cultivated with difficulty on artificial nutrient media, others can lead a saprophytic mode of life and can be satisfied with a negligible collection of organic substances. The latter group enables us to distinguish, in addition to zoonoses, anthroponoses, and anthropozoonoses, still another group of diseases, namely sapronoses (Terskikh, 1958). Of the number of diseases caused by microbes, botulism and also, probably, proteism and melioidosis, can be referred to sapronoses (Fournier, Chambon, 1958).

The genus *Pasteurella* is an interesting example of a gradual increase in the demand for organic substances (Berkman, 1942).

The requirement for sources of energy (and nutrition) varied not only in different species of pathogenic microbes but also in individual strains of one and the same species of microbe (Kuzin, 1946; Stefenson, 1951; Domaradskiy, 1955, 1958; Domaradskiy, Ivanov, 1957).

Attempts to establish a relation between the pathogenicity or virulence of a microbe and the characteristics of its metabolism yields contradictory results. Possibly this is because the concept of "virulence" is still far from sufficiently refined.

We find the most distinct definition of virulence in the work of L. A. Zil'per (1958) who considers that "virulence in contrast to pathogenicity is not a species characteristic. It is a quality inherent to a given strain of microbes, it is an individual characteristic and not a species characteristic. We could say that virulence

is the quality or degree of pathogenicity in a given strain of an infectious agent."

From this position if we are concerned with microbes whose main characteristic as pathogenic agents is their ability to cause toxemia and not an infection proper (*Bac. botulinus*, *Bac. tetani*, *Bac. perfringens* etc.), we can place an equality between virulence and toxigenicity. In addition, we perceive a difference between toxigenicity and toxicity. Toxigenicity is the property of the pathogenicity of a definite group of microbes producing toxins which are easily excreted to the environment in a soluble form and have a characteristic pharmacologic activity. Toxicity is inherent to most bacteria and is associated with decomposition products and cell metabolism. Even cells of nonpathogenic species of microorganisms can have toxicity. This explains in particular the absence of correlation between virulence and "toxigenicity" of the plague microbe (Englesberg et al., 1954) and the multiple rise of the toxicity of its decomposition products when treating a culture with glycocoll (Korobkova, 1957) or surface-active substances (Pannell, 1955; Goodner et al., 1955).

The ineffectiveness or contradictions of the attempts to establish a relation between virulence and the characteristics of metabolism can be demonstrated by the following examples.

As is known Huddleson (1942) and Merz (1938) were the first to show that virulent strains of brucellae has a higher catalase activity than avirulent strains. Later these observations were confirmed by a number of authors, including Schierholz and Jeder (1956). However, according to the data of Guerra (1959), the catalase reaction cannot be used either for differentiation of types of brucellae or for determining the degree of their pathogenicity. Geissler (1959a) arrived at the same conclusion.

The determination of urease activity can serve only to distinguish *Brucella suis* from *Brucella abortus* but is not suitable for determining the degree of pathogenicity of brucellae (Geissler, 1959b).

According to Rockenmacher (1949) a direct relationship exists between the activity of catalase and the virulence of the plague microbe. However, M. N. Dzhabaridze (1953) denies this. It was also established that the virulence of the plague microbe is not associated with fibrinolysin, hyaluronidase, the capacity to coagulate plasma of various species of animals, the oxidation under aerobic and anaerobic conditions of a number of substrates, including amino acids purine and pyrimidine bases, the activity of transaminase, aldolase, phosphatase and a number of other enzymes (Sagar et al., 1956a, b; Saxena et al., 1957; Srikantan et al., 1957a, b; 1958; Misra et al., 1960).

The capacity of strains of *Salmonella typhi* to oxidize glucose tyrosine, and glutamic acid does not depend on the presence in them of  $V_1$ -antigen (Baron, 1955; Costa, Villela, 1959). Judging by the data of Galayev and team (1961) the correlation between pathogenicity and decarboxylase activity of *Escherichia coli* cannot be considered as proved.

Edwards (1960) could not establish a relationship between toxin formation and the activity of catalase for the use of maltose in cultures of *Corynebacterium diphtheriae*.

Fleming and Foshay (1955, 1956) made an interesting attempt to reveal the biochemical nature of the virulence of tularemia bacteria. Judging by the data which can presently be considered only as preliminary, virulent strains of tularemia microbes in contrast to avirulent strains have the capacity to split citrullin and a higher activity than enzymes oxidizing glutamic acid. Conversely, the avirulent strains are distinguished by high activity of the system

aspartate-alanine-transaminase. The oxidation rate of serine or amination of pyruvate is the same for both groups of strains.

On the other hand, in individual cases there is such an expressed dependence between pathogenicity and some aspect of microbial metabolism that the degree of its pathogenicity (i.e., virulence as meant by Zil'ber or toxigenicity) can be judged only on the base of results of in vitro experiments without resorting to experiments on animals. Unfortunately, such a dependence is far from always the exclusive cause of microbial pathogenicity.

In recent years titration methods for  $\alpha$ -toxin of *Bac. perfringens* (or antitoxin) have been widely introduced in practice; these methods following Nagler's principle are based on the decomposition by toxin of the lecithin walls of serum fat particles or of egg lecithovitellin (Nagler, 1941; Heyningen, 1941; Tsekhnovitser et al., 1943; Anan'yeva, 1942; Dotsenko, 1948 and others). However, more accurate methods are those methods in which the lecithin activity of the toxin is dependent by the amount of phosphorus of the split-off phosphorylcholine or by the amount of carbon dioxide displaced by the product of lecithin splitting.

To distinguish pathogenic strains of mycobacteria from nonpathogenic, it is especially important to determine dehydrogenase and catalase activity.

In the presence of glucose, nonpathogenic strains of mycobacteria bleach methylene blue for 1-3 min, whereas strains of avirulent (attenuated) *Mycobacterium tuberculosis* or BCG, for 3-15 min. In contrast to this, suspensions of young virulent *Mycobacterium tuberculosis* do not bleach methylene blue during the course of an hour (Bloch, 1950).

Virulent strains of mycobacteria in comparison with avirulent strains and saprophytic mycobacteria have a higher catalase activity. Inactivation of the catalase of virulent strains occurs at 64-68° and of avirulent strains at 58-60° (Schweiger et al., 1958).

Presently the determination of catalase activity is most frequently used when studying the virulence of drug-resistant strains.

Viallier and Cayré (1958), who established a dissimilar sensitivity to hydrogen peroxide of strains of mycobacteria having similar catalase activity, propose to determine the degree of pathogenicity by the survival rate of cells of these strains in contact with 0.15% hydrogen peroxide for one hour.

Other data on the relation between the enzymic activity of mycobacteria and their virulence can be found in the works of Model' (1958), Segal and Bloch (1956), Kolsut (1959), and others.

A no less striking example is the relation between the characteristics of metabolism and the pathogenicity of staphylococci.

The main criterion for distinguishing in vitro pathogenic strains of staphylococci from nonpathogenic is coagulase (Vygodchikov, 1959; Munch-Petersen, 1961).

In addition to coagulase, phosphatase is an index of staphylococcal pathogenicity (Marinelli et al., 1958). Tests for phosphatase and coagulase coincided in 98.7% of the cases when checking 209 strains of *Staphylococcus aureus* (170 coagulase-positive and 39 coagulase-negative). However upon multiple subculturing the strains sometimes lose phosphatase activity (Pérez-Miravete, Gallardo, 1959).

It was also established that the pathogenicity of staphylococci in the overwhelming majority is correlated with the presence of arginase in it. For instance, out of 70 pathogenic strains, 69 contained arginase, and out of 25 strains considered nonpathogenic only 4 had

arginase (Soru et al., 1957). In addition, the ability only after 2-3 hr growth to form extracellular deoxyribonuclease is characteristic of pathogenic strains of staphylococci; strains of nonpathogenic species of staphylococci also form this enzyme but it can be detected only at considerably later periods, after one or two days (Redaelli, Rosaschino, 1958).

Another test with ammonium molybdenate was proposed for determining the degree of staphylococcal pathogenicity (Myers, 1959).

#### 4.

Each new work confirms the thought first expressed about 100 years ago by Roux: "Death in infectious diseases occurs as a result of intoxication." Therefore it is just for this reason that it is so important to know why under certain conditions the products of the vital activity of microbes are toxic for an organism and how the toxins act. Dubo (1948) said, "Unfortunately, our ignorance in these problems is truly frightening." Possibly Dubo somewhat exaggerated, but there is unconditionally a portion of truth in his words. Our data, which in no way pretends to be complete, show how far we still are from solving these problems.

The main task now is to study the factors causing the pathogenicity of bacteria. At the same time we must stop attempting to identify all toxins with enzymes and devote more attention to the "nonspecific" pathogenetic role of the latter.

#### REFERENCES

- B. V. Anan'yeva, Titrovaniye syvorotok antiperfringes metodom Neglera. Zh. mikrobiol., epidemiol., i immunol., no. 10, 79, 1942.  
G. V. Vyrodchikov, Biorbiologiya i immunologiya stafylokokkovykh zabolevaniy. Medgiz, 1950.

- Ye. L. Gubarev, A. P. Tsyfir'eva, Soderzhaniye aminiaka i glutamina v mozge pri bakterial'nykh intoksikatsiyakh. Vopr. med. khim., 4, 365, 1958.
- Yu. V. Galayev, Dekarpoksilazy aminokislot disenteriynykh mikrobov. Konf. fil. Yuga RSFSR Vses. ob-va fiziologov, biokhimikov i farmakologov. Tez. dokl. P. 62, 1956.
- Yu. V. Galayev, Ye. L. Gubarev, V. P. Davydov, Dekarboksilazy aminokislot u razlichnykh stamov Bact. coli v svyazi s ikh patogennost'yu. V mezhdunar. biokhim. kongr. Referaty sektiionnykh soobshcheniy, 2, 19, 1961.
- Ye. L. Gubarev, Bakteriokhimiya. Gosmedizdat USSR, 1956. Fermenty bakteriy kak patogeneticheskiye faktory pri infektsionnykh zabolevaniyakh. Konf. fil. Yuga RSFSR Vses. ob-va fiziologov, biokhimikov i farmakologov. Tez. dokl. P. 81.
- Ye. L. Gubarev, N. N. Ivanovskiy, Biokhimiya chumnogo mikroba. Medgiz, 1958.
- M. N. Dzhabaridze, Katalaznaya i peroksidaznaya aktivnost' chumnogo i psevdotuberkuleznogo mikrobov. Dissertatsiya. Biblioteka in-ta "Mikrob." Saratov, 1953.
- I. V. Demaradskiy, Aminokislotnyy obmen mikrobov chumy i psevdotuberkuleza. Disseratsiya. Biblioteka inst. "Mikrob." Saratov, 1958. Obshchaya kharakteristika obmena veshchestva chumnogo mikroba. Izv. Irkutskogo protivochumnogo inst. Sibiri i Dal'n Vost., 18, 155, 1961. Ob antifermentakh-antitelakh. Uspekhi sovrem. biol., Vol. 51, No. 4.
- I. V. Demaradskiy, V. A. Ivanov, Nekotoriye dannyye po kul'tivirovaniyu chumnogo mikrobakh na sinteticheskikh sredakh. Zh. mikrobiol., epidemiol., i immunol., No. 2, 54, 1957.
- I. V. Demaradskiy, I. E. Alimova, I. G. Perevalova, Vliyaniye toksina chumnogo mikroba na fosfornyy obmen v pecheni. Vopr. med. khimii, 7, 145, 1961.
- I. V. Demaradskiy, G. A. Yaromyuk, Z. I. Vasil'eva, K voprosy ob otnoshenii chumnogo mikroba k gialuronovoy kislote i nalichii u nego faktora pesproustraneniya. Zh. mikrobiol., epidemiol. i immunol., No. 7, 113, 1960.
- P. S. Lotsenko, Sravnitel'nyy, metod titratsii syvrotki antiperfringes. Ob. nauchn. tr. Irkutskogo inst. epidemiol., 4, 138, 1948.
- G. I. Zepletina, K izucheniyu fermentov prirody toksina chumnogo mikroba, Trudy Rostov-n/a n.-i protivochum. inst., 15, 127, 1959.
- L. A. Sil'ber, Osnovy immunologii, 1958.
- S. Ya. Kaplan'skiy, Khimicheskaya priroda toksinov Bact. perfringens i ikh vliyaniye na protsessy obmena v organisme zhivotnykh i cheloveka. Uspekhi biol. khimii, 2, 292, 1954.
- A. F. Konikov, Nekotoryye zadachi immunokhimicheskogo izucheniya patologii infektsionnogo protsessa. Tr. V sess. AN SSSR, Medgiz. P. 120, 1950.
- Ye. I. Korobkova, Leystviye glikokola na chumnoy mikrob. Soobshch. III. O prirode chumnogo toksina i yego znachenii v aktivnoy immunizatsii. Zh. mikrobiol., epidemiol., i immunol., No. 11, 143, 1957.



- A. P. Kuzin, Khimiya i biokhimiya patogennykh mikrobov. Medgiz, 1946.
- A. I. Mateyev, Botulizm. Medgiz, 1959. Epidemiologiya i profilaktika stolbnyaka, Medgiz, 1960.
- Ye. K. Mishustin, M. I. Pertsovskaya, Mikroorganizmy i samoochishcheniye pochvy, Medgiz, 1954.
- L. M. Model', Biologiya tuberkuleznykh mikobakteriy i immunobiologiya tuberkuleza, Medgiz, 1958.
- T. S. Pashina, Vliyaniya proteoliticheskikh fermentov i nepolnykh (chastichnykh) gidrolizatorov fibrogena na pronitsayemost' kapillyarov. Vopr. med. khimii, 1, 44, 1955.
- Ye. A. Petrosyan, V. P. Zvenigorodskaya, Izucheniye Vi-antigena B. coli i S. typhimur. Zh. mikrobiol., epidemiol. i immunol., No. 10, 114, 1957.
- A. M. Peverenny, Nekotorye predstavleniya o patogeneticheskom znachenii bakterial'nykh proteinaz (k mekhanizmu bakterial'nogo vozdeystviya na kollagenovyye struktury), sb. tr. po biokhim. i fiziol. mikrobov. Gor'kiy, 1961.
- L. M. Fustovalova, Aminokislotnyy sostav kholernykh i kholeropodobnykh vibriionov. Vopr. med. khimii, 7, 265, 1961.
- G. P. Samarina, L. G. Kritsman, L. M. Yakobson, A. Konikova, Aminokislotnyy sostav bakteriy. Biokhimiya, 15, 287, 1950.
- V. I. Tersikh, Sapronozy (O boleznyakh lyudey i zhivotnykh vyzyvayemykh mikrobnymi sposobnymi razmnozheniya vne organizma vo vneshney srede, yavlyayushcheyssya dlya nikh mestom obitaniya). Zh. mikrobiol., epidemiol. i immunol., No. 8, 118, 1958.
- M. M. Tsekhnovitsker, B. L. Shtutina, F. M. Mitel'man, O titratsii in vitro protivogangrenoznykh syvorotok. Zh. mikrobiol., epidemiol. i immunol., No. 6, 48, 1943.
- G. N. Chistovich, O mekhanizme deystviya bakterial'nykh toksinov. Uspekhi sovrem. biol., Vol. 31, 101, 1951.
- V. K. Shaposhnikov, Fiziologiya obmena veshchestv. mikroorganizmov v svyazi s evolyutsiyey funktsiy. Izd. AN SSSR, 1960.
- V. G. Shpikiter, G. A. Levnikova, V. N. Grekhovich, A. I. Solov'yev, Mekhanizme deystviya i nekotorykh svoystvakh kollagenazy. V mezhdunar. biokhim. kongr. Referaty sektsionnykh sobshcheniy. Vol. II, P. 83, 1961.
- Almon L. 1943. The significance of the Vi-antigen. Bacteriol. Rev., 7, 43.
- Avery O., Heidelberger M., Goebel W. 1925. The soluble specific substance of Friedländer's bacillus. II. Chemical and immunological relationships of pneumococcus type II and of a strain of Friedländer's bacillus. J. Expt. Med., 42, 709.
- Baron L. 1955. Enzymic activities of Vi strains of Salmonella typhi and their W variants. Experientia, 11, 268.
- Bennett E., Peterson G., Williams R. 1959. Penicillin sensitivity as compared with nucleic acid phosphorus content of virulent and avirulent strains of Bacillus anthracis and Bacillus cereus. Antibiot. and Chemotherapy, 9, 115.
- Berkman S. 1942. Accessory growth factor requirements of the members of the genus Pasteurella. J. Infect. Diseases, 71, 201.
- Blobel H., Berman D. 1961. Further studies on the in vivo activity of staphylocoagulase. J. Infect. Diseases, 108, 63.
- Blobel H., Berman D., Simon J. 1960. Purification of staphylococcal coagulase. J. Bacteriol., 79, 807.
- Bloch H. 1950. Studies on the virulence of tubercule bacilli. J. Exptl. Med., 91, 2. Cited from L. M. Model', 1958.
- Boake W. 1950. Antistaphylocoagulase in experimental staphylococcal infections. J. Immunol., 70, 89.
- Braude A., Siemieniaski J. 1960. Role of bacterial urease in experimental pyelonephritis. J. Bacteriol., 80, 171.

- Buchler H., Schantz E., Lamanna C. 1947. The elemental and amino acid composition of crystalline *Clostridium botulinum* type A toxin. *J. Biol. Chem.*, 169, 295.  
Cited from K. I. Matveyev, 1959.
- Campbell D., Fourt L. 1939. Immunochemistry of catalase. *J. Biol. Chem.*, 129, 385.
- Carlson A., Kellner A., Bernheimer A. 1956. Selective inhibition by preparations of streptococcal filtrates of the oxidative metabolism of mitochondria procured from rabbit myocardium. *J. Exptl. Med.*, 104, 577.
- Carlson A., Kellner A., Bernheimer A., Freeman E. 1957. A streptococcal enzyme that acts specifically upon diphosphopyridine nucleotide: characterization of the enzyme and its separation from streptolysin. *O. J. Exptl. Med.*, 106, 15.
- Chu H. 1939. The lecithinase of *Bacillus cereus* and its comparison with *Clostridium welchii*  $\alpha$ -toxin. *J. Gen. Microbiol.*, 3, 255.
- Costa G., Villela G. 1959. Oxidation of L-glutamic acid in relation to antigenic composition of *Salmonella typhi*. *Proc. Soc. Exptl. Biol. and Med.*, 100, 714.
- Costlow R. 1958. Lecithinase from *Bacillus anthracis*. *J. Bacteriol.*, 76, 317.
- De launay A., Lebrun J. 1947. Le mode d'action des endotoxines bacteriennes. I. Introduction a une etude generale. *Ann. Inst. Pasteur*, 73, 554.
- Drummond M., Tager M. 1959a. Enzymatic activity of staphylocoagulase. I. Characterization of an esterase associated with purified preparations. *J. Bacteriol.*, 78, 167.— 1959b. Enzymatic activity of staphylocoagulase. II. Dissociation of plasma clotting from tributyrinase activity. *J. Bacteriol.*, 78, 413.
- Jubas, R. 1948. *Bacterial Cell*. IL-1957. **Biochemical Factors in bacterial diseases. IL.**
- Duthie E., Lorenz L. 1952. Staphylococcal coagulase: mode of action and antigenicity. *J. Gen. Microbiol.*, 7, 320.
- Edwards D. 1960. The growth and toxin production of *Corynebacterium diphtheriae* in submerged culture. *J. Gen. Microbiol.*, 22, 658.
- Eisler D. 1961. Coagulation of human plasma by *Pasteurella pestis*. *J. Bacteriol.*, 81, 241.
- Englesberg E., Chen T., Levy J., Foster L., Meyer K. 1954. Virulence in *Pasteurella pestis*. *Science*, 119, 413.
- Fleming D., Foshay L. 1955. Studies on the physiology of virulence of *Pasteurella tularensis*. I. Citrulline ureidase and deamidase activity. *J. Bacteriol.*, 70, 345.— 1956. Studies on the physiology of virulence of *Pasteurella tularensis*. II. Serine deaminase and transaminase activity. *J. Bacteriol.*, 71, 324.
- Fournier J., Chambon L. 1958. Le melioidose et le bacille de Whitmore. Paris.
- Fusillo M., Jaffurs W. 1953. Relationship of coagulase and urease in *Micrococcus pyogenes*. *J. Bacteriol.*, 70, 481.
- Geissler A. 1959a. Die Beziehungen zwischen Katalaseaktivität und Virulenz der Brucella-Bakterien. I. Experimentelle untersuchungen an weissen Mäusen. *Zbl. Veterinärmed.*, 6, 429.— 1959b. Die Bedeutung der Ureaseaktivität der Brucella-Bakterien. *Zbl. Veterinärmed.*, 6, 565.
- Goodner K., Pannell L., Bartell P., Rothstein E. 1953. Toxic end products from *P. pestis*. I. A comparison of lysate toxin with that obtained from the action of bile salts. *J. Infect. Diseases*, 93, 82.
- Guerra M. 1959. Ulteriori ricerche sull'attività catalasica delle brucelle. Nota II. *Nuovi ann. igiene e microbiol.*, 10, 266. Cited from *Biochem. Abs. J.*, No. 7, 9151, 1960.
- Haas R. 1940. Über die Antigeneigenschaften von Papainpräparaten. *Biochem. Z.*, 303, 280.
- Harvey D., Carne Phillida. 1960. Studies on some chemical aspects of the pathological activities of strains of *Escherichia coli* of bovine origin. *J. Compar. Pathol. and Therap.*, 70, 81. Cited from *Biochem. Abs. J.*, 8C595, 1951.
- Hendley Ch., Robbins K., Deetinger B., Clements G., Ercoli N. 1956. Studies on proteolytic enzymes. I. Toxicology of crystalline trypsin and chymotrypsin. *Arch. internat. pharmacodyn.*, 106, 164.
- Heyningen W. van 1911. The biochemistry of the gas gangrene toxins. I. Estimation of the  $\alpha$ -toxin of *Cl. welchii* type A. *Biochem. J.*, 35, 1246.
- Heyningen W. van 1960. Bacterial toxins. Oxford.
- Heyningen, W. Toxic proteins. In the Book: "Proteins," edited by H. Neurath and K. Bayley, IL, 3, part I, P. 639.
- Holtmann D., Peterson W. 1952. Evidence of a staphylococcal neurotoxin. *J. Bacteriol.*, 64, 590.
- Howell S. 1932. Antiurease formation in the hen. *Proc. Soc. Exptl. Biol. and Med.*, 29, 759.
- Howes E., Mandl I., Zaffuto S. 1960. Toxicity of enzymes from *Clostridium histolyticum*. *J. Bacteriol.*, 79, 191.
- Huddleson I. 1942. *Univ. Mich. Agri. Exp. Station. Techn. Bull.*, 182, January. Cited from Rockenmacher M., 1949.
- Iseki Shoji, Furukawa Ken. 1959. On blood group specific decomposing enzymes derived from bacteria. *Proc. Japan Acad.*, 35, 620. Cited from *Biochem. Abs. J.*, No. 5, 558, 1961.
- Isibasi K. 1959. Nippon ika daigaku zasshi. *J. Nippon Med. School.*, 26, 195. Cited from *Biochem. Abs. J.*, No. 9, 12252, 1960.

- Jawetz E., Meyer K. 1944. Studies on plague immunity in experimental animals. II. Some factors of the immunity mechanism in bubonic plague. *J. Immunol.*, 49, 15.
- Jenkin C., Rowley D. 1959. Possible factors in the pathogenesis of cholera. *Brit. J. Exptl Pathol.*, 40, 474.
- Kellner A., Robertson T. 1951a. Selective necrosis of cardiac and skeletal muscle induced experimentally by means of proteolytic enzyme solutions given intravenously. *J. Exptl Med.*, 99, 387.—1951b. Cardiac lesions produced experimentally in animals given crystalline streptococcal proteinase intravenously. *Amer. J. Pathol.*, 30, 636.
- Keppie J., Smith H., Harris-Smith P. 1955. The chemical basis of the virulence of *Bacillus anthracis*. III. The role of the terminal bacteriaemia in death of guinea-pigs from anthrax. *Brit. J. Exptl Pathol.*, 36, 315.
- Kirk J. 1933. The concentration of soy bean urease. A new method for the purification of enzymes. *J. Biol. Chem.*, 100, 637.
- Kirk J., Sumner J. 1931. Anturease. *J. Biol. Chem.*, 94, 21.—1934. The reaction between crystalline urease and anturease. *J. Immunol.*, 26, 495.
- Kleeberg S. 1958. Harufush. *J. Med. Assoc. Israel*, 54, 201.
- Kolsnt H. 1959. Aktywnosc enzymow i zjadliwosc pratkow gruźlicy opornych na hydrazyd kwasu izonikotynowego. *Gruźlica*, 27, 277.
- Lie E. 1960. Über die Fermentniture der Staphylokoagulase. *Arch. Hyg. und Bakteriol.*, 144, 32.
- Lise, n., 1958. **Biochemistry of autotrophic bacteria. II.**
- Luippold G. 1942. Typhoid vaccine studies. V. Studies on the relationship between the antigenic content and the immunogenic properties of bacterial suspensions. *Amer. J. Hyg.*, 36, 354.—1944. Antityphoid activity of V<sub>1</sub>-antigen from extragenic sources. *Science*, 99, 497.
- Macfarlane M., Knight J. 1941. The biochemistry of bacterial toxins. I. The lecithinase activity of *Cl. welchii* toxins. *Biochem. J.*, 35, 884.
- Macfarlane R., MacLennan J. 1945. The toxemia of gas gangrene. *Lancet*, 11, 328.
- Macfarlane R., Oakley C., Anderson C. 1941. Haemolysis and the production of opalescence in serum and lecithovitellin by the  $\alpha$ -toxin of *Clostridium welchii*. *J. Pathol. Bact.*, 52, 59.
- Marinelli M., Tonelli E., Venturi R. 1958. Attività fosfatasi e potere patogeno dello stafilococco. *Giorn. batteriol. virol. ed immunol.*, 51, 566.
- Merz P. 1938. Über die Katalasen der Brucellen. Inaugural Dissertation for the degree of Doctor of Veterinary Medicine. Universität of Zürich. Lhr. no. Rockenmacher M., 1949.
- Miles E., Miles A. 1950. The relation of toxicity and enzyme activity in the lecithinases of *Clostridium bifementans* and *Clostridium welchii*. *J. Gen. Microbiol.*, 4, 22.
- Misra J., Agarwala S., Shrivastava D. 1960. Study in the Enzyme Make-Up of *Pasteurella pestis*. Part VII. Deamination of Purine and Pyrimidine compounds by virulent and avirulent strains. *Ind. J. Med. Res.*, 48, 1.
- Munch-Petersen E. 1961. Staphylococcal coagulase. A survey of contributions during the last two decades. *Zbl. Bakteriol. Parasitenkunde, Infektionskrankh. und Hyg.*, 178, 377.
- Myers D. 1959. Observations on a chemical test related to the virulence of *Staphylococcus aureus*. *Amer. J. Clin. Pathol.*, 31, 128.
- Nagler F. 1941. The estimation of the combining power of *Cl. welchii* (type A) toxoid. *J. Pathol. Bact.*, 52, 105.
- Neill J. 1939. Immunological relationships between pneumococci and various species of fungi. 3rd. Internat. Congr. Microbiol., p. 807.
- Oakley C., Warrack G., Heyning W. 1946. The collagenase ( $K$ -toxin) of *Cl. welchii*, type A. *J. Pathol. Bact.*, 58, 229.
- Oakley C., Warrack G., Warren M. 1948. Kappa ( $\kappa$ )- and lambda ( $\lambda$ )-antigens of *Cl. welchii*. *J. Pathol. Bact.*, 60, 495.
- Pannell L. 1955. The effect of surface active agents of the toxicity of *P. pestis* lysates. *J. Infect. Diseases*, 96, 54.
- Pappenheimer A. 1947a. Diphtheria toxin. Effect of iron on toxin and porphyrin production. *J. Biol. Chem.*, 167, 251.—1947b. Bacterial toxins. *Federat. Proc.*, 6, 479.
- Pappenheimer A., Hendee E. 1947. Diphtheria toxin. IV. The iron enzymes of *Corynebacterium diphtheriae* and their possible relation to diphtheria toxin. *J. Biol. Chem.*, 171, 701.
- Pappenheimer A., Williams C. 1952. The effects of diphtheria toxin on the *Cecropia silkworm*. *J. Gen. Physiol.*, 35, 727.
- Pérez-Miravete A., Gallardo R. 1959. La prueba de la fosfatasa comparada con otras pruebas de patogenicidad de *Staphylococcus aureus*. *Ciencia (Mexico)*, 19, 29. Cited from *Biochem. Abs. J.*, No. 5, 6100, 1960.
- Phear E., Ruebner B. 1956. The in vitro production of ammonium and amines by intestinal bacteria in relation to nitrogen toxicity as a factor in hepatic coma. *Brit. J. Exptl Pathol.*, 37, 253.
- Ramon G. 1957. Ferments, anaferments, antiferments étude immunologique. *Rev. immunol. et de therapie antimicrobienne*, 21, 221.

- Redaelli G., Rosaschino F. 1958. Recherche sulla desossiribonucleasi stafilococcali. Nuovi ann. ingiene e microbiol., 9, 545. Cited from Biochem. Abs. J., No. 18, 239-9, 1959.
- Richou R., Chirol C., Durieux R., Quinchon C. 1960. Sur le passage, à travers le placenta, de principes capables de neutraliser certaines propriétés enzymatiques des toxines staphylococciques. Compt. rend. Acad. sci., 251, 3136.
- Richou R., Quinchon C., Chirol. 1961. Sur la présence, dans les  $\gamma$ -globulines d'origine humaine, de principes capables de neutraliser certaines propriétés enzymatiques des toxines staphylococciques. Compt. rend. Acad. sci., 252, 226.
- Robb-Smith A. 1945. Tissue changes induced by Cl. welchii type A filtrates. Lancet, II, 362.
- Robinson J., Thatcher F., Montford J. 1960. Studies with Staphylococcal toxin. V. Possible identification of alpha hemolysin with a proteolytic enzyme. Canad. J. Microbiol., 6, 183.
- Rocha e Silva M. 1939a. Action of trypsin on the smooth muscle of isolated guinea-pig intestine. Compt. rend. Soc. biol., 130, 181.—1939b. Action of hyppsin on the guinea-pig uterus. C. R. Soc. Biol., 130, 184.—1940. Pharmacology of trypsin II. Action on blood pressure in cat, dog and rabbit. Arch. Exptl. Pathol. Pharmacol., 194, 351.
- Rockenmacher M. 1949. Relationship of catalase activity to virulence in Pasteurella pestis. Proc. Soc. Exptl. Biol. and Med., 71, 99.
- Sagar P., Agarwala S., Shrivastava D. 1956a. Studies in the Enzyme Make-Up of Pasteurella pestis. Part I. Deamination of amino acids by virulent and avirulent strains. Ind. J. Med. Res., 44, 385.—1956b. Studies in the Enzyme Make-Up of Pasteurella pestis. Part II. Phosphatase activity of virulent and avirulent strains. Ind. J. Med. Res., 44, No. 4.
- Sasaki T. 1958. Japan J. Bacteriol., 13, 105. Cited from Biochem. Abs. J., No. 23, 20153, 1958.
- Saxena K., Sagar P., Agarwala S., Shrivastava D. 1957. Studies in the Enzyme Make-Up of Pasteurella pestis. Part IV. Transamination reactions of virulent and avirulent strains. Ind. J. Med. Res., 45, No. 2.
- Schierholz H., Jäder R. 1956. Kinetische untersuchungen über den durch Bangbakterien verursachten peroxylzerfall. Zbl. Bakteriol. Parasitenkunde, Infektionskrankh. u. Hyg., Abt. I. Orig., 167, 281.
- Schweiger O., Roman E., Soos I. 1958. Studies on the catalase enzyme of mycobacteria I. Correlation between catalase activity and virulence of mycobacteria under certain experimental conditions. Amer. Rev. Tuberc., 77, 146.
- Segal W., Bloch H. 1956. Biochemical differentiation of Mycobacterium tuberculosis grown in vivo and in vitro. J. Bacteriol., 72, 132.
- Smith D., Johnstone J. 1958. Brit. J. Exptl. Path., 34, 165. Cited from H. Blobel and D. Berman, 1961.
- Soru E., Nestorescu N., Popovici M., Ionescu M. 1957. Un non test de différentiere a stafilococilor patogeni, testul arginazic. Studii și cercetari inframicrobiol., microbiol și parazitol., 8, 395. Cited from Biochem. Abs. J., No. 2, 1373, 1959.
- Srikantan T., Agarwala S., Shrivastava D. 1957a. Studies in the Enzyme Make-Up of P. pestis. Part III. Oxidative metabolism of virulent and avirulent strains. Ind. J. Med. Res., 45, No. 2.—1957b. Studies in the Enzyme Make-Up of P. pestis. Part V. Dehydrogenase activity of virulent and avirulent strains. Ind. J. Med. Res., 45, 467.
- Srikantan T., Krishna Murti C., Shrivastava D. 1958. Studies in the Enzyme Make-Up of P. pestis. Part VI. Aldolase activity of virulent and avirulent strains. Ind. J. Med. Res., 46, 1.
- Stevenson, M. 1951. Metabolism of bacteria.
- Strauss N., Hendee E. 1959. The effect of diphtheria toxin on the metabolism of HeLa cells. J. Exptl. Med., 109, 145.
- Stuart C., Wheeler K., Rustigian R., Zimmermann A. 1943. Biochemical and antigenic relationships of the paracolon bacteria. J. Bacteriol., 45, 101.
- Sugiyama H., Dack G. 1955. Apyrase in partially purified staphylococcal enterotoxin. J. Infect. Diseases, 96, 286.
- Takatori, 1955. Nihon densen be gakkai dzassi, 29, 67. Cited from Biochem. Abs. J., No. 5, 4723, 1957.
- Thompson T., Militzer W., Giorgi C. 1958. Partial denaturation of a bacterial aldolase without loss of activity. J. Bacteriol., 76, 337.
- Todd E., Lancefield R. 1928. Variants of hemolytic streptococci; their relation to type-specific substance, virulence and toxin. J. Exptl. Med., 48, 751.
- Tracey M. 1954. Principles of Biochemistry. Pitman, London. Cited after Lise, 1958.
- Waller J., Cayré R. 1958. Activité enzymatiques et virulence de souches de Mycobacterium tuberculosis isoniazidorésistantes. Compt. rend. Soc. biol., 152, 1156.
- Welch, H. G. 1929. Chemistry of Immunity. II.
- Woolley, D. A study of antimetabolites. II, 1954.
- Zamecnik P., Lipmann F. 1947. A study of the competition of lecithin and antitoxin for Cl. welchii lecithinase. J. Exptl. Med., 85, 395.
- Zeller E., Märitz A. 1944. Helv. chim. acta, 27, 1888. Cited after D. Somner and G. Somner. Chemistry of enzymes and methods of their investigation. II, 1948.

# DISTRIBUTION LIST

DEPARTMENT OF DEFENSE	Nr. Copies	MAJOR AIR COMMANDS	Nr. Copies
		AFSC	
		SCFDD	1
		ASTIA	25
HEADQUARTERS USAF		TDBTL	5
		TDBDP	5
AFCIN-3D2	1	TDEWA (Bullard)	1
ARL (ARB)	1	SSD (SSF)	2
AFCIN-M	1	AMD (AMRF)	1
		ESD (ESY)	2
OTHER AGENCIES			
CIA	1		
NSA	6		
DIA	9		
AID	2		
OTS	2		
AEC	2		
PWS	1		
NASA	1		
ARMY	3		
NAVY	3		
NAFEC	1		
PGE	12		
RAND	1		